



Journal of Multivariate Analysis 98 (2007) 774–788

www.elsevier.com/locate/jmva

A multilinear form inequality

Frederic Picard

Department of Mathematics, Indiana University, Bloomington, IN 47405, USA

Received 23 June 2005

Available online 19 January 2006

Abstract

An inequality of Interpolation type for Multilinear Forms with a two-part dependence condition is proved. It generalizes the work of Bradley and Bryc [Theorem 3.6, Multilinear forms and measures of dependence between random variables, *J. Multivariate Anal.* 16 (1985) 335–367] and Prakasa Rao [Bounds for r th order joint cumulant under r th order strong mixing, *Statist. Probab. Lett.* 43 (1999) 427–431].

© 2005 Elsevier Inc. All rights reserved.

AMS 2000 subject classification: 60E15

Keywords: Interpolation; Multilinear forms; Cumulant; Two-part dependence condition

1. Introduction

We start with some definitions and notations. If W is nonnegative random variable on a probability space (Ω, \mathcal{F}, P) , then for each $z \in (0, 1)$, define the quantile

$$Q_W(z) := \inf\{t \geq 0 : P(W > t) \leq z\}.$$

We notice that $t \geq Q_W(z)$ if and only if $P(W > t) \leq z$.

For any set A , I_A or $I(A)$ will denote the indicator function of A . Let \mathcal{A} be any σ -field $\subset \mathcal{F}$. A simple random variable is a random variable for which the range is a finite set. We denote by $\mathcal{S}(\mathcal{A})$ the set of \mathbb{C} -valued \mathcal{A} -measurable simple random variables. If $\mathcal{A}_1, \dots, \mathcal{A}_r$ are σ -fields, a function $F : \mathcal{S}(\mathcal{A}_1) \times \dots \times \mathcal{S}(\mathcal{A}_r) \rightarrow \mathbb{C}$ is said to be a multilinear form if it is linear in each of

E-mail address: frpicard@indiana.edu.

its variables. If in addition for all $X_1 \in \mathcal{S}(\mathcal{A}_1), \dots, X_r \in \mathcal{S}(\mathcal{A}_r)$ we have that

$$|F(X_1, \dots, X_r)| \leq \|X_k\|_1 \cdot \prod_{i \neq k} \|X_i\|_\infty \quad \text{for } k = 1, \dots, r \quad (1)$$

then we say that F is a multilinear product form. Multilinear forms arise in the study of dependence coefficients. A very common example of a multilinear form is $\text{Cov}(X_1, X_2)$. More generally when $r \geq 2$, there is the multivariate cumulant:

$$\text{Cum}(X_1, \dots, X_r) := \sum (-1)^{p-1} (p-1)! \left(E \prod_{j \in v_1} X_j \right) \dots \left(E \prod_{j \in v_p} X_j \right),$$

where the sum is taken over the partitions (v_1, \dots, v_p) of $\{1, \dots, r\}$ for $p = 1, \dots, r$. If one multiplies the cumulant by $(\sum_{p=1}^r p^r \cdot (p-1)!)^{-1}$, then it satisfies (1) and thus it becomes a product form. In [3], some inequalities concerning multilinear product forms are proved using interpolation theory. The main goal of this paper is to prove the following theorem:

Theorem 1. Suppose r is a positive integer and that $\mathbf{p} = (p_1, \dots, p_r) \in [1, \infty]^r$ is such that $\sum_{i=1}^r p_i^{-1} \leq 1$. Define the number $c = \text{Card}\{i \in \{1, \dots, r\} : p_i < \infty\} - \sum_{i=1}^r p_i^{-1}$. Then there exists a constant $C = C(\mathbf{p})$ that is a function only of \mathbf{p} such that the following holds: suppose (Ω, \mathcal{F}, P) is a probability space, $\mathcal{A}_1, \dots, \mathcal{A}_r$ are σ -subfields of \mathcal{F} , $F : \mathcal{S}(\mathcal{A}_1) \times \dots \times \mathcal{S}(\mathcal{A}_r) \rightarrow \mathbb{C}$ is a multilinear product form and α and λ are numbers in $[0, 1]$ such that for every choice of events $A_i \in \mathcal{A}_i$, one has that

$$|F(I_{A_1}, \dots, I_{A_r})| \leq \alpha + \lambda \cdot \prod_{i=1}^r [P(A_i)]^{1/p_i}; \quad (2)$$

then for all $X_1 \in \mathcal{S}(\mathcal{A}_1), \dots, X_r \in \mathcal{S}(\mathcal{A}_r)$, we have

$$|F(X_1, \dots, X_r)| \leq 4^r \cdot \int_0^\alpha \prod_{i=1}^r \mathcal{Q}_{|X_i|}(z) dz + C \cdot \lambda (1 - \log(\lambda))^c \cdot \prod_{i=1}^r \|X_i\|_{p_i}.$$

The case when $\alpha = 0$ was proved in [3]. Rio proved several results related to the case $\lambda = 0$ (see for example [8, Theorem 1.1]). In [6] Prakasa Rao studied the case in which F is the cumulant function and $\sum_{i=1}^r p_i^{-1} = 1$. By a simple application of Riesz–Thorin’s Multilinear Interpolation Theorem (see [1, p. 18, Exercise 13]), if F is a multilinear product form and $\mathbf{p} = (p_1, \dots, p_r) \in [1, \infty]^r$ is such that $\sum_{i=1}^r p_i^{-1} \leq 1$, then for all $X_1 \in \mathcal{S}(\mathcal{A}_1), \dots, X_r \in \mathcal{S}(\mathcal{A}_r)$, we have $F(X_1, \dots, X_r) \leq \prod_{i=1}^r \|X_i\|_{p_i}$ (see [3, p. 349] for details). However, Theorem 1 gives a bound for F that gets small as λ and α get small.

The assumption (2) is called a two-part dependence condition. Peligrad [5] proved a CLT for a class of strictly stationary sequences that satisfy a two-part dependence condition. These sequences arise, for example, when one applies a certain nonlinear smoothing algorithm of Tukey [9] (the “3R” or “running median” smoother) to strictly stationary ρ -mixing sequences (see [4]).

The CLT in [5] involved the parameters λ and α both getting small. In such problem as finding a CLT for the estimators of cumulants (themselves or their spectral densities), we would like the parameters λ and α to be small so that Theorem 1 could be applied.

2. Preliminaries

We will use the following lemmas.

Lemma 1. Suppose that the simple random variables $X_1 \in \mathcal{S}(\mathcal{A}_1), \dots, X_r \in \mathcal{S}(\mathcal{A}_r)$ are non-negative real-valued. Then

$$F(X_1, \dots, X_r) = \int_0^\infty \cdots \int_0^\infty F(I(X_1 > t_1), \dots, I(X_r > t_r)) dt_1 \dots dt_r.$$

Proof. Let $0 = b_{i,0} < b_{i,1} < b_{i,2} < \cdots < b_{i,n_i}$ the values in the range of $X_i, i = 1, \dots, r$ with the possible exception of $b_{i,0} = 0$. Then

$$X_i = \sum_{j=0}^{n_i} b_{i,j} \cdot I(X_i = b_{i,j}) = \sum_{j=1}^{n_i} (b_{i,j} - b_{i,j-1}) I(X_i \geq b_{i,j}).$$

Using the multilinearity of F we get

$$\begin{aligned} F(X_1, \dots, X_r) &= \sum_{j_1=1}^{n_1} \cdots \sum_{j_r=1}^{n_r} \prod_{i=1}^r (b_{i,j_i} - b_{i,(j_i-1)}) F(I(X_1 \geq b_{1,j_1}), \dots, I(X_r \geq b_{r,j_r})) \\ &= \sum_{j_1=1}^{n_1} \cdots \sum_{j_r=1}^{n_r} \int_{b_{1,(j_1-1)}}^{b_{1,j_1}} \cdots \int_{b_{r,(j_r-1)}}^{b_{r,j_r}} F(I(X_1 > t_1), \dots, I(X_r > t_r)) dt_1 \dots dt_r \\ &= \int_0^\infty \cdots \int_0^\infty F(I(X_1 > t_1), \dots, I(X_r > t_r)) dt_1 \dots dt_r. \quad \square \end{aligned}$$

The following result is essentially due to Rio [7].

Lemma 2. Suppose that X_1, \dots, X_r are nonnegative real-valued simple random variables. Then

$$\int_0^\infty \cdots \int_0^\infty \min\{P(X_1 > t_1), \dots, P(X_r > t_r), \alpha\} dt_1 \dots dt_r = \int_0^\alpha \prod_{i=1}^r Q_{X_i}(z) dz.$$

Proof. A proof is given by Prakasa Rao [6, Eq. (1.1)]. \square

Lemma 3. Suppose L and M are each a positive integer, and $t > 0$. Then

$$\sum_{n_1=L}^\infty \cdots \sum_{n_M=L}^\infty 2^{-t \max\{n_1, \dots, n_M\}} \leq M! \frac{2^{-t \cdot L}}{(1 - 2^{-t})^M}.$$

Proof. We notice that

$$\begin{aligned} & \sum_{n_1=L}^{\infty} \cdots \sum_{n_M=L}^{\infty} 2^{-t \max\{n_1, \dots, n_M\}} \\ & \leq M! \sum_{n_1=L}^{\infty} \sum_{n_2=n_1}^{\infty} \cdots \sum_{n_{M-1}=n_{M-2}}^{\infty} \sum_{n_M=n_{M-1}}^{\infty} 2^{-t \cdot n_M} \end{aligned}$$

and the result follows by applying M times the geometric series formula. \square

We recall the fact that any complex random variable Y can be written as $Y = Y_1 - Y_2 + \sqrt{-1} \cdot (Y_3 - Y_4)$, where Y_j for $j = 1, \dots, 4$ are nonnegative random variables with $Y_j \leq |Y|$. First recall that any complex random variable can be written in the form $Y = \Re(Y) + \sqrt{-1} \cdot \Im(Y)$, where $\Re(Y)$ and $\Im(Y)$ denote respectively the real and imaginary part of Y . Then the random variables $Y_1 = \max\{\Re(Y), 0\}$, $Y_2 = \max\{-\Re(Y), 0\}$, $Y_3 = \max\{\Im(Y), 0\}$ and $Y_4 = \max\{-\Im(Y), 0\}$ will have the desired property. This, the multilinearity of F and the facts that $\mathcal{Q}_{Y_j} \leq \mathcal{Q}_{|Y|}$, $j = 1, \dots, 4$ show that in order to prove Theorem 1, it is enough to prove the following proposition.

Proposition 1. *Suppose the hypothesis of Theorem 1 holds, and in addition the random variables X_1, \dots, X_r are real-valued and nonnegative. Then*

$$|F(X_1, \dots, X_r)| \leq \int_0^\alpha \prod_{i=1}^r \mathcal{Q}_{X_i}(z) dz + \frac{C}{4^r} \cdot \lambda(1 - \log(\lambda))^c \prod_{i=1}^r \|X_i\|_{p_i}.$$

It follows from Proposition 1 that Theorem 1 remains valid with its final inequality replaced by the one above if we have the extra assumption that the random variables X_1, \dots, X_r are real and nonnegative.

The proof of Proposition 1 begins as follows:

Since F is a product form, for each $k \in \{1, \dots, r\}$, we have

$$\begin{aligned} & |F(I(X_1 > t_1), \dots, I(X_r > t_r))| \\ & \leq \|I(X_k > t_k)\|_1 \prod_{i \neq k} \|I(X_i > t_i)\|_\infty \\ & \leq \|I(X_k > t_k)\|_1 \\ & = P(X_k > t_k). \end{aligned} \tag{3}$$

Using the main assumption (2),

$$|F(I(X_1 > t_1), \dots, I(X_r > t_r))| \leq \alpha + \lambda \cdot \prod_{i=1}^r [P(X_i > t_i)]^{1/p_i}. \tag{4}$$

Combining (3) and (4) we obtain

$$\begin{aligned} & |F(I(X_1 > t_1), \dots, I(X_r > t_r))| \\ & \leq \min \left\{ P(X_1 > t_1), \dots, P(X_r > t_r), \alpha + \lambda \cdot \prod_{i=1}^r [P(X_i > t_i)]^{1/p_i} \right\} \end{aligned}$$

$$\leq \min \{P(X_1 > t_1), \dots, P(X_r > t_r), \alpha\} \\ + \min \left\{ P(X_1 > t_1), \dots, P(X_r > t_r), \lambda \cdot \prod_{i=1}^r [P(X_i > t_i)]^{1/p_i} \right\}.$$

Using this inequality and Lemma 1 we obtain

$$|F(X_1, \dots, X_r)| \\ = \left| \int_0^\infty \dots \int_0^\infty F(I(X_1 > t_1), \dots, I(X_r > t_r)) dt_1 \dots dt_r \right| \\ \leq \int_0^\infty \dots \int_0^\infty |F(I(X_1 > t_1), \dots, I(X_r > t_r))| dt_1 \dots dt_r \\ \leq \int_0^\infty \dots \int_0^\infty \min \{P(X_1 > t_1), \dots, P(X_r > t_r), \alpha\} dt_1 \dots dt_r \\ + \int_0^\infty \dots \int_0^\infty \min \left\{ P(X_1 > t_1), \dots, P(X_r > t_r), \right. \\ \left. \lambda \cdot \prod_{i=1}^r [P(X_i > t_i)]^{1/p_i} \right\} dt_1 \dots dt_r$$

and by Lemma 2 we know that

$$\int_0^\infty \dots \int_0^\infty \min \{P(X_1 > t_1), \dots, P(X_r > t_r), \alpha\} dt_1 \dots dt_r = \int_0^\alpha \prod_{i=1}^r \mathcal{Q}_{X_i}(z) dz.$$

So it is enough to prove the following:

$$\int_0^\infty \dots \int_0^\infty \min \left\{ P(X_1 > t_1), \dots, P(X_r > t_r), \lambda \prod_{i=1}^r [P(X_i > t_i)]^{1/p_i} \right\} dt_1 \dots dt_r \\ \leq \frac{C}{4^r} \cdot \lambda (1 - \log(\lambda))^c \cdot \prod_{i=1}^r \|X_i\|_{p_i}.$$

So we see that the proof of Proposition 1 reduces to the following lemma:

Lemma 4. Suppose r is a positive integer and $\mathbf{p} = (p_1, \dots, p_r) \in [1, \infty]^r$ with $\sum_{i=1}^r p_i^{-1} \leq 1$, $\lambda \in [0, 1]$. Define the number $c = c(\mathbf{p}) := \text{Card}\{k : p_k < \infty\} - \sum_{i=1}^r p_i^{-1}$. Then there exists a constant $C = C(\mathbf{p})$ that is a function only of \mathbf{p} such that the following holds:

If X_1, \dots, X_r are nonnegative simple random variables, then we have

$$\int_0^\infty \dots \int_0^\infty \min \left\{ P(X_1 > t_1), \dots, P(X_r > t_r), \lambda \cdot \prod_{i=1}^r [P(X_i > t_i)]^{1/p_i} \right\} dt_1 \dots dt_r \\ \leq C \cdot \lambda (1 - \log(\lambda))^c \cdot \prod_{i=1}^r \|X_i\|_{p_i}.$$

(Notice that C is basically the same as in Proposition 1 and Theorem 1 with the difference that it has been multiplied by $(4^r)^{-1}$.) Here and for the rest of the paper, \log will denote the logarithm to the base e . As usual, \log_2 denotes the logarithm to the base 2.

3. Proof of Lemma 4

If $\lambda = 0$, the proof is trivial. So we can assume $0 < \lambda \leq 1$. We will separate the proof into four cases.

Case 1: $\mathbf{p} \in (1, \infty)^r$ and $\sum_{i=1}^r p_i^{-1} < 1$.

The proof will be written out here under the extra assumption that $r \geq 2$. If instead $r = 1$, the proof is similar but much simpler. For each $j \in \{1, 2, \dots, r\}$, define the positive constant $K_j = K_j(\mathbf{p})$ (where $\mathbf{p} = (p_1, \dots, p_r)$) as follows:

$$\begin{aligned} K_j &= K_j(\mathbf{p}) \\ &= 2 \left(\prod_{k=1}^r \frac{1}{2^{1/p_k} - 1} \right) \\ &\quad \times \sum \left[(r - (1 + \text{Card } A))! \cdot \frac{1}{\left(1 - 2^{-(1/p_j + 1 - \sum_{k=1}^r 1/p_k)}\right)^{r - (1 + \text{Card } A)}} \right], \end{aligned} \quad (5)$$

where the sum is taken over all sets $A \subseteq \{2, 3, \dots, r\}$ (including the empty set). Note that it can be rewritten

$$\begin{aligned} K_j &= K_j(\mathbf{p}) \\ &= 2 \left(\prod_{k=1}^r \frac{1}{2^{1/p_k} - 1} \right) \\ &\quad \times \sum_{l=0}^{r-2} \binom{r-1}{l} \left[(r - (1 + l))! \cdot \frac{1}{\left(1 - 2^{-(1/p_j + 1 - \sum_{k=1}^r 1/p_k)}\right)^{r-1-l}} \right]. \end{aligned}$$

For each $j \in \{1, \dots, r\}$, define the positive constant $C_j = C_j(\mathbf{p})$ as follows:

$$\begin{aligned} C_j &= C_j(\mathbf{p}) \\ &= 2^r K_j + 2^{r+2} \cdot [\max\{p_1, p_2, \dots, p_r\}]^{r-1}. \end{aligned}$$

Define the positive $C = C(\mathbf{p})$ as follows:

$$C = \sum_{j=1}^r C_j.$$

Now suppose X_1, \dots, X_r are nonnegative simple random variables. Let $H : [0, \infty)^r \rightarrow \mathbb{R}$ be defined as

$$H(t_1, \dots, t_r) := \min \left\{ P(X_1 > t_1), \dots, P(X_r > t_r), \lambda \cdot \prod_{i=1}^r [P(X_i > t_i)]^{1/p_i} \right\}. \quad (6)$$

Our aim is to show that

$$\int_0^\infty \dots \int_0^\infty H(t_1, \dots, t_r) dt_1 \dots dt_r \leq C \cdot \lambda (1 - \log(\lambda))^C \cdot \prod_{i=1}^r \|X_i\|_{p_i}.$$

For each $i = 1, \dots, r$, define the nondecreasing sequence $a_{i,n}$ by

$$\begin{aligned} a_{i,0} &= 0, \\ a_{i,n} &= \mathcal{Q}_{X_i}(2^{-n}) \text{ for } n \geq 1. \end{aligned}$$

Also we define $J_{i,n} := [a_{i,n}, a_{i,n+1})$. If $a_{i,n} = a_{i,n+1}$ then $J_{i,n} = \emptyset$. We see that

$$2^{-n-1} < P(X_i > x_i) \leq 2^{-n}, x_i \in J_{i,n}, i = 1, \dots, r.$$

In [2], it is proved that for $k = 1, \dots, r$

$$\frac{1}{2} \sum_{n=0}^{\infty} 2^{-n} |J_{k,n}|^{p_k} \leq EX_k^{p_k}$$

which implies

$$\left(\sum_{n=0}^{\infty} 2^{-n} |J_{k,n}|^{p_k} \right)^{1/p_k} \leq 2^{1/p_k} \|X_k\|_{p_k}. \quad (7)$$

We see that

$$\{J_{1,n_1} \times \dots \times J_{r,n_r} : n_1, \dots, n_r \in \{0, 1, 2, \dots\}\}$$

is a collection of pairwise disjoint subsets of $[0, \infty)^r$ such that the support H is contained in the union of that collection. Therefore if we define for each $i_1, \dots, i_r \in \{0, 1, 2, 3, \dots\}$,

$$G_{i_1, \dots, i_r} := \int \dots \int_{(x_1, \dots, x_r) \in J_{1,i_1} \times \dots \times J_{r,i_r}} H(x_1, \dots, x_r) dx_1 \dots dx_r,$$

it is easy to see that

$$\int_0^\infty \dots \int_0^\infty H(t_1, \dots, t_r) dt_1 \dots dt_r = \sum_{i_1=0}^\infty \dots \sum_{i_r=0}^\infty G_{i_1, \dots, i_r}.$$

Since each of the G_{i_1, \dots, i_r} 's is nonnegative, we have that

$$\begin{aligned} & \sum_{i_1=0}^\infty \dots \sum_{i_r=0}^\infty G_{i_1, \dots, i_r} \\ & \leq \sum_{i_1=0}^\infty \sum_{i_2=0}^\infty \dots \sum_{i_r=0}^\infty G_{i_1, i_1+i_2, \dots, i_1+i_r} + \sum_{i_1=0}^\infty \sum_{i_2=0}^\infty \dots \sum_{i_r=0}^\infty G_{i_1+i_2, i_1, i_1+i_3, \dots, i_1+i_r} \\ & \quad + \dots + \sum_{i_1=0}^\infty \sum_{i_2=0}^\infty \dots \sum_{i_r=0}^\infty G_{i_1+i_2, i_2+i_3, \dots, i_1+i_r, i_1}. \end{aligned} \quad (8)$$

We will find an upper bound for

$$\sum_{i_1=0}^\infty \dots \sum_{i_r=0}^\infty G_{i_1, i_1+i_2, \dots, i_1+i_r} = \sum_{i_2=0}^\infty \dots \sum_{i_r=0}^\infty \left(\sum_{i_1=0}^\infty G_{i_1, i_1+i_2, \dots, i_1+i_r} \right). \quad (9)$$

The upper bounds for the other terms of (8) can be found in a similar way. In order to find the upper bound for (9), we will find two bounds for

$$\sum_{i_1=0}^{\infty} G_{i_1, i_1+i_2, \dots, i_1+i_r}. \quad (10)$$

The first bound will be used when all the integers i_2, \dots, i_r are “reasonably small” and the second one will be used when at least one of the integers i_2, \dots, i_r is “large”.

By (6), for $(x_1, \dots, x_r) \in J_{1,i_1} \times \dots \times J_{r,i_r}$ we have

$$H(x_1, \dots, x_r) \leq \min\{2^{-i_1}, \dots, 2^{-i_r}, \lambda \cdot 2^{-i_1/p_1 - \dots - i_r/p_r}\}$$

and hence

$$G_{i_1, \dots, i_r} \leq |J_{1,i_1}| \dots |J_{r,i_r}| \min\{2^{-i_1}, \dots, 2^{-i_r}, \lambda \cdot 2^{-i_1/p_1 - \dots - i_r/p_r}\}. \quad (11)$$

Let s be the number such that

$$\frac{1}{p_1} + \frac{1}{p_2} + \dots + \frac{1}{p_r} + \frac{1}{s} = 1.$$

Obviously, $s > 1$. We defined s in order to apply Hölder’s inequality. Choose M the nonnegative integer such that

$$M \leq s \cdot \log_2(1/\lambda) < M + 1.$$

For any nonnegative integers i_2, \dots, i_r ,

$$\begin{aligned} & \sum_{i_1=0}^{\infty} G_{i_1, i_1+i_2, \dots, i_1+i_r} \\ &= \sum_{i_1=0}^{M-1} G_{i_1, i_1+i_2, \dots, i_1+i_r} + \sum_{i_1=M}^{\infty} G_{i_1, i_1+i_2, \dots, i_1+i_r}. \end{aligned} \quad (12)$$

First we will get a bound for the first term in the right-hand side of (12). Using (11), Hölder’s inequality (with exponents p_1, p_2, \dots, p_r and s), (7) and the definition of M we get

$$\begin{aligned} & \sum_{i_1=0}^{M-1} G_{i_1, i_1+i_2, \dots, i_1+i_r} \\ & \leq \sum_{i_1=0}^{M-1} \lambda \cdot 2^{-i_1/p_1} \cdot 2^{-(i_1+i_2)/p_2} \dots 2^{-(i_1+i_r)/p_r} \cdot |J_{1,i_1}| \cdot |J_{2,i_1+i_2}| \dots |J_{r,i_1+i_r}| \\ &= \lambda \cdot \sum_{i_1=0}^{M-1} |J_{1,i_1}| \cdot 2^{-(i_1)/p_1} \left\{ \prod_{k=2}^r |J_{k,i_1+i_k}| \cdot 2^{-(i_1+i_k)/p_k} \right\} \cdot 1 \\ & \leq \lambda \left(\sum_{i_1=0}^{M-1} |J_{1,i_1}|^{p_1} \cdot 2^{-i_1} \right)^{1/p_1} \end{aligned}$$

$$\begin{aligned}
& \times \left\{ \prod_{k=2}^r \left(\sum_{i_1=0}^{M-1} |J_{k,i_1+i_k}|^{p_k} \cdot 2^{-(i_1+i_k)} \right)^{1/p_k} \right\} \left(\sum_{m=0}^{M-1} 1^s \right)^{1/s} \\
& \leq \lambda (2 \cdot E|X_1|^{p_1})^{1/p_1} \cdot \left\{ \prod_{k=2}^r (2 \cdot E|X_k|^{p_k})^{1/p_k} \right\} M^{1/s} \\
& = \lambda \cdot 2^{1/p_1+\dots+1/p_r} \cdot M^{1/s} \prod_{k=1}^r \|X_k\|_{p_k} \\
& \leq \lambda \cdot 2 \cdot (s \cdot \log_2(1/\lambda))^{1/s} \prod_{k=1}^r \|X_k\|_{p_k} \\
& \leq 4 \cdot \lambda \cdot (\log_2(1/\lambda))^{1/s} \prod_{k=1}^r \|X_k\|_{p_k}.
\end{aligned}$$

For the last inequality, we used the fact that $s^{1/s} < 2$ which is easy to prove using calculus.

Now, we get an upper bound for the second term in the right-hand side of (12). Let $i_{\max} := \max\{i_2, \dots, i_r\}$. Notice that $2^{-i_{\max}+i_2/p_2+\dots+i_r/p_r} \leq 1$. Thus using (11), Hölder's inequality, (7) and the definition of M we have,

$$\begin{aligned}
& \sum_{i_1=M}^{\infty} G_{i_1,i_1+i_2,\dots,i_1+i_r} \\
& \leq \sum_{i_1=M}^{\infty} |J_{1,i_1}| \cdot |J_{2,i_1+i_2}| \cdot \dots \cdot |J_{r,i_1+i_r}| \cdot 2^{-i_1-i_{\max}} \\
& = 2^{-i_{\max}+i_2/p_2+\dots+i_r/p_r} \cdot \sum_{i_1=M}^{\infty} |J_{1,i_1}| \cdot 2^{-i_1/p_1} \left\{ \prod_{k=2}^r |J_{k,i_1+i_k}| \cdot 2^{-(i_1+i_k)/p_k} \right\} \cdot 2^{-i_1/s} \\
& \leq \sum_{i_1=M}^{\infty} |J_{1,i_1}| \cdot 2^{-i_1/p_1} \left\{ \prod_{k=2}^r |J_{k,i_1+i_k}| \cdot 2^{-(i_1+i_k)/p_k} \right\} \cdot 2^{-i_1/s} \\
& \leq \left(\sum_{i_1=M}^{\infty} |J_{1,i_1}|^{p_1} 2^{-i_1} \right)^{1/p_1} \left\{ \prod_{k=2}^r \left(\sum_{i_1=M}^{\infty} |J_{k,i_1+i_k}|^{p_k} \cdot 2^{-(i_1+i_k)} \right)^{1/p_k} \right\} \\
& \quad \times \left(\sum_{i_1=M}^{\infty} 2^{-i_1} \right)^{1/s} \\
& \leq 2^{1/p_1} \cdot \|X_1\|_{p_1} \left\{ \prod_{k=2}^r 2^{1/p_k} \|X_k\|_{p_k} \right\} (2^{-M+1})^{1/s} \\
& = 2^{1/p_1+\dots+1/p_r} \cdot \left\{ \prod_{k=1}^r \|X_k\|_{p_k} \right\} \cdot (2^{-M+1})^{1/s} \\
& \leq 2^{1/p_1+\dots+1/p_r} \cdot \left\{ \prod_{k=1}^r \|X_k\|_{p_k} \right\} \cdot (2^{-s \log_2(1/\lambda)+2})^{1/s}
\end{aligned}$$

$$\begin{aligned}
&= 2^{1/p_1+\dots+1/p_r} \cdot \left\{ \prod_{k=1}^r \|X_k\|_{p_k} \right\} \cdot \lambda \cdot 2^{2/s} \\
&= 2 \cdot 2^{1/s} \cdot \lambda \prod_{k=1}^r \|X_k\|_{p_k} \\
&\leq 4 \cdot \lambda \cdot \prod_{k=1}^r \|X_k\|_{p_k}.
\end{aligned}$$

So combining the last two inequalities in (12), we get that for any nonnegative integers i_2, \dots, i_r ,

$$\sum_{i_1=0}^{\infty} G_{i_1, i_1+i_2, \dots, i_1+i_r} \leq 4\lambda \cdot (1 + (\log_2(1/\lambda))^{1/s}) \prod_{k=1}^r \|X_k\|_{p_k}. \quad (13)$$

Now we get another inequality for (10). We recall that we defined $i_{\max} = \max\{i_2, \dots, i_r\}$. By (11), Hölder's inequality and (7)

$$\begin{aligned}
&\sum_{i_1=0}^{\infty} G_{i_1, i_1+i_2, \dots, i_1+i_r} \\
&\leq \sum_{i_1=0}^{\infty} |J_{1, i_1}| \cdot |J_{2, i_1+i_2}| \dots |J_{r, i_1+i_r}| \cdot 2^{-(i_1+i_{\max})} \\
&= 2^{-i_{\max}+i_2/p_2+\dots+i_r/p_r} \sum_{i_1=0}^{\infty} |J_{1, i_1}| \cdot 2^{-i_1/p_1} \left\{ \prod_{k=2}^r |J_{k, i_1+i_k}| 2^{-(i_1+i_k)/p_k} \right\} \cdot 2^{-i_1/s} \\
&\leq 2^{-i_{\max}+i_2/p_2+\dots+i_r/p_r} \\
&\quad \times \left(\sum_{i_1=0}^{\infty} |J_{1, i_1}|^{p_1} 2^{-i_1} \right)^{1/p_1} \cdot \left\{ \prod_{k=2}^r \left(\sum_{i_1=0}^{\infty} |J_{k, i_1+i_k}|^{p_k} 2^{-(i_1+i_k)} \right)^{1/p_k} \right\} \cdot \left(\sum_{i_1=0}^{\infty} 2^{-i_1} \right)^{1/s} \\
&\leq 2^{-i_{\max}+i_2/p_2+\dots+i_r/p_r} \cdot 2^{1/p_1} \|X_1\|_{p_1} \dots 2^{1/p_r} \|X_r\|_{p_r} \cdot 2^{1/s} \\
&= 2 \cdot 2^{-i_{\max}+i_2/p_2+\dots+i_r/p_r} \cdot \prod_{i=1}^r \|X_i\|_{p_i}. \quad (14)
\end{aligned}$$

We are now ready to find an upper bound for (9). Let L be the nonnegative integer such that

$$L - 1 \leq \max \left\{ \frac{p_1 \cdot s}{s + p_1}, \dots, \frac{p_r \cdot s}{s + p_r} \right\} \cdot \log_2 \left(\frac{1}{\lambda} \right) < L.$$

Then

$$\begin{aligned}
&\sum_{i_1=0}^{\infty} \dots \sum_{i_r=0}^{\infty} G_{i_1, i_1+i_2, \dots, i_1+i_r} \\
&= \sum_{i_2=1}^L \dots \sum_{i_r=1}^L \sum_{i_1=1}^{\infty} G_{i_1, i_1+i_2, \dots, i_1+i_r} + \sum_{i_{\max} \geq L} \sum_{i_1=0}^{\infty} G_{i_1, i_1+i_2, \dots, i_1+i_r} \quad (15)
\end{aligned}$$

(here $\sum_{i_{\max} \geq L}$ means the sum over all $(i_2, \dots, i_r) \in \{0, 1, 2, \dots\}^{r-1}$ such that $i_{\max} \geq L$).

We will later bound the first term in the right-hand side of (15) using (13). Now we turn our attention to the second term. Since by (14) we have

$$\begin{aligned} \sum_{i_{\max} \geq L} \sum_{i_1=0}^{\infty} G_{i_1, i_1+i_2, \dots, i_1+i_r} \\ \leq 2 \cdot \prod_{i=1}^r \|X_i\|_{p_i} \cdot \sum_{i_{\max} \geq L} 2^{-i_{\max}+i_2/p_2+\dots+i_r/p_r}, \end{aligned}$$

we will get an upper bound for that last sum. For that purpose, note that

$$\begin{aligned} \sum_{i_{\max} \geq L} 2^{-i_{\max}+i_2/p_2+\dots+i_r/p_r} \\ = \sum_A \left(\sum_{i_{\sigma(2)}=0}^{L-1} \dots \sum_{i_{\sigma(a)}=0}^{L-1} \sum_{i_{\sigma(a+1)}=L}^{\infty} \dots \sum_{i_{\sigma(r)}=L}^{\infty} 2^{-i_{\max}+i_{\sigma(2)}/p_{\sigma(2)}+\dots+i_{\sigma(r)}/p_{\sigma(r)}} \right) \\ = \sum_A \left(\sum_{i_{\sigma(2)}=0}^{L-1} \dots \sum_{i_{\sigma(a)}=0}^{L-1} 2^{i_{\sigma(2)}/p_{\sigma(2)}+\dots+i_{\sigma(a)}/p_{\sigma(a)}} \right) \\ \times \left(\sum_{i_{\sigma(a+1)}=L}^{\infty} \dots \sum_{i_{\sigma(r)}=L}^{\infty} 2^{-i_{\max}+i_{\sigma(a+1)}/p_{\sigma(a+1)}+\dots+i_{\sigma(r)}/p_{\sigma(r)}} \right), \end{aligned} \quad (16)$$

where each sum is taken over all subsets $A \subsetneq \{2, 3, \dots, r\}$ and for each A , we choose one permutation $\sigma : \{2, 3, \dots, r\} \rightarrow \{2, 3, \dots, r\}$ such that $\sigma(2), \dots, \sigma(a) \in A$ and $\sigma(a+1), \dots, \sigma(r) \notin A$ (where $a = 1 + \text{Card}(A)$). Using lemma 3 we get that

$$\begin{aligned} \sum_{i_{\sigma(a+1)}=L}^{\infty} \dots \sum_{i_{\sigma(r)}=L}^{\infty} 2^{-i_{\max}+i_{\sigma(a+1)}/p_{\sigma(a+1)}+\dots+i_{\sigma(r)}/p_{\sigma(r)}} \\ \leq \sum_{i_{\sigma(a+1)}=L}^{\infty} \dots \sum_{i_{\sigma(r)}=L}^{\infty} 2^{-i_{\max} \cdot (1/p_1+1/p_{\sigma(2)}+\dots+1/p_{\sigma(a)}+1/s)} \\ \leq (r-a)! \cdot \frac{2^{-L \cdot (1/p_1+1/p_{\sigma(2)}+\dots+1/p_{\sigma(a)}+1/s)}}{(1-2^{-(1/p_1+1/s)})^{r-a}} \end{aligned} \quad (17)$$

and by a direct computation using the geometric series formula

$$\begin{aligned} \sum_{i_{\sigma(2)}=0}^{L-1} \dots \sum_{i_{\sigma(a)}=0}^{L-1} 2^{i_{\sigma(2)}/p_{\sigma(2)}+\dots+i_{\sigma(a)}/p_{\sigma(a)}} \\ = \frac{2^{L/p_{\sigma(2)}} - 1}{2^{1/p_{\sigma(2)}} - 1} \dots \frac{2^{L/p_{\sigma(a)}} - 1}{2^{1/p_{\sigma(a)}} - 1} \\ \leq \frac{2^{L/p_{\sigma(2)}}}{2^{1/p_{\sigma(2)}} - 1} \dots \frac{2^{L/p_{\sigma(a)}}}{2^{1/p_{\sigma(a)}} - 1} \\ = \frac{2^{L \cdot (1/p_{\sigma(2)}+\dots+1/p_{\sigma(a)})}}{(2^{1/p_{\sigma(2)}} - 1) \dots (2^{1/p_{\sigma(a)}} - 1)} \\ \leq 2^{L \cdot (1/p_{\sigma(2)}+\dots+1/p_{\sigma(a)})} \left(\prod_{i=1}^r \frac{1}{2^{1/p_i} - 1} \right). \end{aligned} \quad (18)$$

So using (17) and (18) into (16)

$$\begin{aligned}
 & \sum_{i_{\max} \geq L} \sum_{i_1=0}^{\infty} G_{i_1, i_1+i_2, \dots, i_1+i_r} \\
 & \leq 2 \cdot \prod_{i=1}^r \|X_i\|_{p_i} \sum_{i_{\max} \geq L} 2^{-i_{\max}+i_2/p_2+\dots+i_r/p_r} \\
 & \leq 2 \cdot \prod_{i=1}^r \|X_i\|_{p_i} \sum_A (r-a)! \cdot \frac{2^{-L \cdot (1/p_1+1/s)}}{(1-2^{-(1/p_1+1/s)})^{r-a}} \prod_{i=1}^r \frac{1}{2^{1/p_i}-1} \\
 & = 2 \cdot \prod_{i=1}^r \|X_i\|_{p_i} 2^{-L \cdot (1/p_1+1/s)} \left(\sum_A (r-a)! \cdot \frac{1}{(1-2^{-(1/p_1+1/s)})^{r-a}} \prod_{i=1}^r \frac{1}{2^{1/p_i}-1} \right) \\
 & = K_1 \cdot 2^{-L \cdot (1/p_1+1/s)} \cdot \prod_{i=1}^r \|X_i\|_{p_i}, \tag{19}
 \end{aligned}$$

where K_1 was defined by Eq. (5). Using (13), (19) and the definition of L

$$\begin{aligned}
 & \sum_{i_1=0}^{\infty} \sum_{i_2=0}^{\infty} \dots \sum_{i_r=0}^{\infty} G_{i_1, i_1+i_2, \dots, i_1+i_r} \\
 & = \sum_{i_2=0}^{\infty} \dots \sum_{i_r=0}^{\infty} \left(\sum_{i_1=0}^{\infty} G_{i_1, i_1+i_2, \dots, i_1+i_r} \right) \\
 & = \sum_{i_2=0}^{L-1} \dots \sum_{i_r=0}^{L-1} \sum_{i_1=0}^{\infty} G_{i_1, i_1+i_2, \dots, i_1+i_r} + \sum_{i_{\max} \geq L} \sum_{i_1=0}^{\infty} G_{i_1, i_1+i_2, \dots, i_1+i_r} \\
 & \leq \sum_{i_2=0}^{L-1} \dots \sum_{i_r=0}^{L-1} 4\lambda \cdot (1 + (\log_2(1/\lambda))^{1/s}) \prod_{k=1}^r \|X_k\|_{p_k} + \sum_{i_{\max} \geq L} \sum_{i_1=0}^{\infty} G_{i_1, i_1+i_2, \dots, i_1+i_r} \\
 & = 4\lambda \cdot L^{r-1} \cdot (1 + (\log_2(1/\lambda))^{1/s}) \prod_{k=1}^r \|X_k\|_{p_k} + \sum_{i_{\max} \geq L} \sum_{i_1=0}^{\infty} G_{i_1, i_1+i_2, \dots, i_1+i_r} \\
 & \leq 4\lambda \cdot L^{r-1} \cdot (1 + (\log_2(1/\lambda))^{1/s}) \prod_{k=1}^r \|X_k\|_{p_k} + K_1 \cdot 2^{-L \cdot (\frac{1}{p_1} + \frac{1}{s})} \prod_{i=1}^r \|X_i\|_{p_i} \\
 & \leq 4\lambda \left(\max \left\{ \frac{p_1 \cdot s}{s + p_1}, \dots, \frac{p_r \cdot s}{s + p_r} \right\} \log_2(1/\lambda) + 1 \right)^{r-1} (1 + (\log_2(1/\lambda))^{1/s}) \prod_{k=1}^r \|X_k\|_{p_k} \\
 & \quad + K_1 \cdot \lambda \cdot \prod_{i=1}^r \|X_i\|_{p_i} \\
 & \leq 4\lambda [\max \{p_1, p_2, \dots, p_r\}]^{r-1} (1 + \log_2(1/\lambda))^{r-1} (1 + (\log_2(1/\lambda))^{1/s}) \prod_{k=1}^r \|X_k\|_{p_k} \\
 & \quad + K_1 \cdot \lambda \cdot \prod_{i=1}^r \|X_i\|_{p_i}.
 \end{aligned}$$

For the next to last inequality, we used the following facts that follow from the definition of L :

$$-L < -\log_2(1/\lambda) \frac{p_1 \cdot s}{s + p_1}$$

and

$$L \leq \max \left\{ \frac{p_1 \cdot s}{s + p_1}, \dots, \frac{p_r \cdot s}{s + p_r} \right\} \cdot \log_2(1/\lambda) + 1.$$

By using the fact that $1 \leq (1 + \log_2(1/\lambda))^{r-1+1/s}$, proving $1 + (\log_2(1/\lambda))^{1/s} \leq 2^{1-1/s} \cdot (1 + \log_2(1/\lambda))^{1/s}$ using Hölder's inequality in two dimensions and other elementary facts, we see that

$$\sum_{i_1=0}^{\infty} \sum_{i_2=0}^{\infty} \cdots \sum_{i_r=0}^{\infty} G_{i_1, i_1+i_2, i_1+i_3, \dots, i_1+i_r} \leq C_1 \cdot \lambda(1 + \log(1/\lambda))^{r-1+1/s} \cdot \prod_{i=1}^r \|X_i\|_{p_i}.$$

Similarly we can prove that

$$\begin{aligned} \sum_{i_1=0}^{\infty} \sum_{i_2=0}^{\infty} \cdots \sum_{i_r=0}^{\infty} G_{i_1+i_2, i_1, i_1+i_3, \dots, i_1+i_r} &\leq C_2 \cdot \lambda(1 + \log(1/\lambda))^{r-1+1/s} \cdot \prod_{i=1}^r \|X_i\|_{p_i} \\ \vdots \\ \sum_{i_1=0}^{\infty} \sum_{i_2=0}^{\infty} \cdots \sum_{i_r=0}^{\infty} G_{i_1+i_2, \dots, i_1+i_r, i_1} &\leq C_r \cdot \lambda(1 + \log(1/\lambda))^{r-1+1/s} \cdot \prod_{i=1}^r \|X_i\|_{p_i}. \end{aligned}$$

Thus,

$$\begin{aligned} &\int_0^{\infty} \cdots \int_0^{\infty} H(t_1, \dots, t_r) dt_1 \dots dt_r \\ &= \sum_{i_1=0}^{\infty} \cdots \sum_{i_r=0}^{\infty} G_{i_1, \dots, i_r} \\ &\leq \sum_{i_1=0}^{\infty} \sum_{i_2=0}^{\infty} \cdots \sum_{i_r=0}^{\infty} G_{i_1, i_1+i_2, \dots, i_1+i_r} + \sum_{i_1=0}^{\infty} \sum_{i_2=0}^{\infty} \cdots \sum_{i_r=0}^{\infty} G_{i_1+i_2, i_1, i_1+i_3, \dots, i_1+i_r} \\ &\quad + \cdots + \sum_{i_1=0}^{\infty} \sum_{i_2=0}^{\infty} \cdots \sum_{i_r=0}^{\infty} G_{i_1+i_2, i_2+i_3, \dots, i_1+i_r, i_1} \\ &\leq C_1 \cdot \lambda(1 + \log(1/\lambda))^{r-1+1/s} \cdot \prod_{i=1}^r \|X_i\|_{p_i} \\ &\quad + C_2 \cdot \lambda(1 + \log(1/\lambda))^{r-1+1/s} \cdot \prod_{i=1}^r \|X_i\|_{p_i} \\ &\quad \vdots \quad \quad \quad \vdots \end{aligned}$$

$$\begin{aligned}
& + C_r \cdot \lambda(1 + \log(1/\lambda))^{r-1+1/s} \cdot \prod_{i=1}^r \|X_i\|_{p_i} \\
& = C \cdot \lambda(1 + \log(1/\lambda))^{r-1+1/s} \cdot \prod_{i=1}^r \|X_i\|_{p_i}
\end{aligned}$$

since $C = C_1 + \dots + C_r$. This completes the proof of the Case 1.

Case 2: $\mathbf{p} \in (1, \infty)^r$ and $\sum_{i=1}^n p_i^{-1} = 1$.

Define $C = C(\mathbf{p})$ as in Case 1. We can choose a sequence $\{\mathbf{p}_n = (p_{1,n}, \dots, p_{r,n})\}$ in $(1, \infty)^r$ such that $\sum_{i=1}^r p_{i,n}^{-1} < 1$ and $\mathbf{p}_n \rightarrow \mathbf{p}$. Thus by Case 1, for each $n \in \{1, 2, \dots\}$

$$\begin{aligned}
& \int_0^\infty \dots \int_0^\infty \min \left\{ P(X_1 > t_1), \dots, P(X_r > t_r), \lambda \cdot \prod_{i=1}^r [P(X_i > t_i)]^{1/p_{i,n}} \right\} dt_1 \dots dt_r \\
& \leq C(\mathbf{p}_n) \cdot \lambda(1 - \log(\lambda))^{c(\mathbf{p}_n)} \cdot \prod_{i=1}^r \|X_i\|_{p_{i,n}}. \tag{20}
\end{aligned}$$

We leave to the reader to check that as $n \rightarrow \infty$, $C(\mathbf{p}_n) \rightarrow C(\mathbf{p})$, $c(\mathbf{p}_n) \rightarrow c(\mathbf{p}) = r - 1$ and that $\|X_i\|_{p_{i,n}} \rightarrow \|X_i\|_{p_i}$ for $i = 1, \dots, r$ (note that the X_i 's are simple). So by taking the limit $n \rightarrow \infty$ in (20) we get the result for Case 2.

Case 3: $\mathbf{p} \in (1, \infty]^r$ and $\sum_{i=1}^n p_i^{-1} \leq 1$.

Now we allow the possibility that some p_i 's are equal to infinity. Without loss of generality, we can assume that for some $m \in \{0, \dots, r\}$, $p_i < \infty$ for $i \leq m$ and $p_i = \infty$ for $i > m$. The proof will be written out here for the case $1 \leq m \leq r - 1$. (If instead $m = 0$, the proof is quite trivial; and the case $m = r$ is covered in Cases 1 and 2.) We define $C = C(\mathbf{p}) := C(p_1, \dots, p_m)$ from Case 1 and 2.

$$\begin{aligned}
& \int_0^\infty \dots \int_0^\infty \min \left\{ P(X_1 > t_1), \dots, P(X_r > t_r), \lambda \cdot \prod_{i=1}^r [P(X_i > t_i)]^{1/p_i} \right\} dt_1 \dots dt_r \\
& = \int_0^\infty \dots \int_0^\infty \min \left\{ P(X_1 > t_1), \dots, P(X_r > t_r), \lambda \cdot \prod_{i=1}^m [P(X_i > t_i)]^{1/p_i} \right\} dt_1 \dots dt_r \\
& \leq \int_{t_1=0}^\infty \dots \int_{t_m=0}^\infty \int_{t_{m+1}=0}^{\|X_{m+1}\|_\infty} \dots \int_{t_r=0}^{\|X_r\|_\infty} \\
& \quad \times \min \left\{ P(X_1 > t_1), \dots, P(X_r > t_r), \lambda \cdot \prod_{i=1}^m [P(X_i > t_i)]^{1/p_i} \right\} dt_1 \dots dt_r \\
& \leq \int_{t_1=0}^\infty \dots \int_{t_m=0}^\infty \int_{t_{m+1}=0}^{\|X_{m+1}\|_\infty} \dots \int_{t_r=0}^{\|X_r\|_\infty} \\
& \quad \times \min \left\{ P(X_1 > t_1), \dots, P(X_k > t_m), \lambda \cdot \prod_{i=1}^m [P(X_i > t_i)]^{1/p_i} \right\} dt_1 \dots dt_r \\
& = \int_0^\infty \dots \int_0^\infty \min \left\{ P(X_1 > t_1), \dots, P(X_k > t_m), \right.
\end{aligned}$$

$$\lambda \cdot \prod_{i=1}^m [P(X_i > t_i)]^{1/p_i} \Bigg\} dt_1 \dots dt_m \prod_{i=m+1}^r \|X_i\|_\infty$$

$$\leq C \cdot \lambda (1 - \log(\lambda))^{m - \sum_{i=1}^r p_i^{-1}} \cdot \prod_{i=1}^r \|X_i\|_{p_i}.$$

For the last inequality we applied the result obtained in Cases 1 and 2. This proves Case 3.

Case 4: $\mathbf{p} \in [1, \infty]^r$.

If $\mathbf{p} \in (1, \infty]$ then we are in one of the previous cases. If one p_i 's is equal to 1, then the condition $\sum_{i=1}^r p_i^{-1} = 1$ implies that all the other p_i 's are equal to ∞ . So we may assume without loss of generality that $p_1 = 1$ and $p_2 = \dots = p_r = \infty$. We define $C = C(\mathbf{p}) = 1$. Then

$$\int_0^\infty \dots \int_0^\infty \min \left\{ P(X_1 > t_1), \dots, P(X_r > t_r), \lambda \cdot \prod_{i=1}^r [P(X_i > t_i)]^{1/p_i} \right\} dt_1 \dots dt_r$$

$$= \int_0^\infty \dots \int_0^\infty \min \{ \lambda P(X_1 > t_1), \dots, P(X_r > t_r) \} dt_1 \dots dt_r$$

$$= \int_{t_1=0}^\infty \int_{t_2=0}^{\|X_2\|_\infty} \dots \int_{t_r=0}^{\|X_r\|_\infty}$$

$$\times \min \{ \lambda P(X_1 > t_1), \dots, P(X_r > t_r) \} dt_1 \dots dt_r$$

$$\leq \int_0^\infty \lambda P(X_1 > t_1) dt_1 \prod_{i=2}^r \|X_i\|_\infty^r = \lambda \|X_1\|_1 \prod_{i=2}^r \|X_i\|_\infty^r.$$

This proves Case 4 and thus Lemma 4 is proved. \square

Acknowledgements

I thank Richard Bradley for having suggested the problem. I also thank the reviewer for the helpful comments and suggestions.

References

- [1] J. Bergh, J. Löfström, *Interpolation Spaces*, Springer, Berlin, 1976.
- [2] R.C. Bradley, A covariance inequality under a two-part dependence assumption, *Statist. Probab. Lett.* 30 (1996) 287–293.
- [3] R.C. Bradley, W. Bryc, Multilinear forms and measures of dependence between random variables, *J. Multivariate Anal.* 16 (1985) 335–367.
- [4] W. Bryc, M. Peligrad, The central limit theorem for Tukey's $3R$ smoother, *Statist. Probab. Lett.* 13 (1992) 29–37.
- [5] M. Peligrad, On the central limit theorem for weakly dependent sequences with a decomposed strong mixing coefficient, *Stochastic Process.* 42 (1992) 181–193.
- [6] B.L.S. Prakasa Rao, Bounds for r th order joint cumulant under r th order strong mixing, *Statist. Probab. Lett.* 43 (1999) 427–431.
- [7] E. Rio, Covariance inequalities for the strongly mixing processes, *Ann. Inst. H. Poincaré Probab. Statist.* 29 (1993) 587–597.
- [8] E. Rio, *Théorie Asymptotique des Processus Aléatoires Faiblement Dépendants*, Mathématiques & Applications, vol. 31, Springer, Berlin, 2000.
- [9] J.W. Tukey, *Exploratory Data Analysis*, Addison-Wesley, Reading, MA, 1977.